



US005795720A

United States Patent [19]**Henco et al.**[11] **Patent Number:** **5,795,720**[45] **Date of Patent:** **Aug. 18, 1998**

[54] **PROCESS AND DEVICE FOR THE SEPARATION AND DETECTION OF COMPONENTS OF A MIXTURE OF MATERIALS BY TEMPERATURE GRADIENT GEL ELECTROPHORESIS**

[75] Inventors: **Karsten Henco, Erkrath; Detlev Riesner; Gerhard Steger**, both of Düsseldorf, all of Germany

[73] Assignee: **Qiagen GmbH, Hilden, Germany**

[21] Appl. No.: **558,594**

[22] Filed: **Oct. 31, 1995**

Related U.S. Application Data

[63] Continuation of Ser. No. 200,551, Feb. 22, 1994, abandoned, which is a continuation of Ser. No. 847,761, filed as PCT/EP90/01366 Aug. 18, 1990, abandoned.

[30] **Foreign Application Priority Data**

Aug. 19, 1989	[DE]	Germany	39 27 467.5
Mar. 6, 1990	[DE]	Germany	40 06 974.5

[51] **Int. Cl.⁶** **C12Q 1/68**

[52] **U.S. Cl.** **435/6; 935/77; 935/78**

[58] **Field of Search** **435/6; 935/77, 935/78, 91.2**

[56] **References Cited****U.S. PATENT DOCUMENTS**

4,671,870	6/1987	Tompa et al.	210/149
5,066,377	11/1991	Rosenbaum et al.	204/182.8

FOREIGN PATENT DOCUMENTS

0198207	3/1986	European Pat. Off.
0318273	5/1989	European Pat. Off.

OTHER PUBLICATIONS

Val C. Sheffield et al., "Attachment of a 40-base pair G+C-rich sequence (GC-clamp) to genomic DNA fragments by the polymerase chain . . ." *Proc. Natl. Acad. Sci., USA*, vol. 86, pp. 232-236, 1989.

Ann-Christine Sylvanen et al., "Quantification of polymerase chain reaction products by affinity-based hybrid collection", *Nucleic Acids Research*, vol. 16, No. 23, pp. 11327-11338, 1988.

Roger M. Wartell et al., "Detecting base pair substitutions in DNA fragments by temperature-gradient gel electrophoresis", *Nucleic Acids Research*, vol. 18, No. 9, pp. 2699-2705, 1990.

Richard M. Myers et al., "Nearly all single base substitutions in DNA fragments joined to a GC-clamp can be detected by . . .", *Nucleic Acids Research*, vol. 13, No. 9, pp. 3131-3145, 1985.

Richard M. Myers et al., "Detection & Localization of Single Base Changes by Denaturing Gradient Gel Electrophoresis", *Methods in Enzymology*, vol. 155, pp. 501-527, 1987.

Detlev Riesner et al., "Temperature-gradient gel electrophoresis of nucleic acids: Analysis of conformational transitions, sequence variations, . . .", *Electrophoresis*, vol. 10 (5-6), pp. 377-389, 1989.

Primary Examiner—Lisa B. Arthur

Attorney, Agent, or Firm—Jacobson, Price, Holman & Stern, PLLC

[57] **ABSTRACT**

The invention relates to a process for the separation and detection of components of a mixture of materials by temperature gradient gel electrophoresis, wherein either

a spatial temperature gradient is built up by spatially separated temperature levels, or

a time temperature gradient, or

a temperature gradient is built up by combination of spatial and timewise temperature gradient.

The temperature levels for building up the spatial temperature gradient are adjusted by controllable heating or cooling devices.

To build up the time temperature gradient, the temperature level at each point of the separation path within the separation medium may be optionally adjusted time-dependently by means of controllable heating or cooling devices. There is described a device for performing the process with controllable heating or cooling devices to build up temperature gradients, a hollow body arranged between the temperature levels which contains the medium used for separation, and a thermostat jacket enclosing the hollow body.

12 Claims, 16 Drawing Sheets